

Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Responsive Neurostimulation for the Treatment of Refractory Focal Epilepsy is addressed separately in medical policy 00674.

Note: Vagus Nerve Stimulation is addressed separately in medical policy 00134.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- Benefits are available in the member's contract/certificate, and
- Medical necessity criteria and guidelines are met.

Based on review of available data, the Company may consider unilateral deep brain stimulation (DBS) of the thalamus in individuals with disabling, medically unresponsive tremor due to essential tremor (ET) or Parkinson's disease (PD) to be **eligible for coverage**.**

Based on review of available data, the Company may consider bilateral deep brain stimulation (DBS) of the thalamus in individuals with disabling, medically unresponsive tremor in both upper limbs due to essential tremor (ET) or Parkinson's disease (PD) to be **eligible for coverage**.**

Patient Selection Criteria

Coverage eligibility will be considered for deep brain stimulation (DBS) of the thalamus when the following criteria are met:

Disabling, medically unresponsive tremor defined as:

- Tremor causing significant limitation in daily activities; and
- Inadequate control by maximal dosage of medication for at least three months before implant.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

Based on review of available data, the Company may consider unilateral or bilateral deep brain stimulation (DBS) of the globus pallidus or subthalamic nucleus when patient selection criteria are met to be **eligible for coverage.****

Patient Selection Criteria

Coverage eligibility will be considered for unilateral or bilateral deep brain stimulation (DBS) of the globus pallidus or subthalamic nucleus when all of the following criteria are met:

- Parkinson disease and ALL of the following:
 - A good response to levodopa; AND
 - Motor complications not controlled by pharmacologic therapy; AND
 - One of the following:
 - A minimum score of 30 points on the motor portion of the Unified Parkinson Disease Rating Scale when the patient has been without medication for approximately 12 hours OR
 - Parkinson disease for at least 4 years
- Individuals older than 7 years with chronic, intractable (drug-refractory) primary dystonia, including generalized and/or segmental dystonia, hemidystonia, and cervical dystonia (torticollis).

When Services Are Considered Not Medically Necessary

Based on review of available data, the Company may consider deep brain stimulation (DBS) to be **not medically necessary**** when the following contraindications are present:

- Individuals who are not good surgical risks because of unstable medical problems or because of the presence of a cardiac pacemaker; or
- Individuals who have medical conditions that require repeated magnetic resonance imaging (MRI); or
- Individuals who have dementia that may interfere with the ability to cooperate; or
- Individuals who have had botulinum toxin injections within the last six months.

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

Based on review of available data, the Company considers deep brain stimulation for other movement disorders, including but not limited to tardive dyskinesia and post-traumatic dyskinesia to be **investigational.***

Based on review of available data, the Company considers deep brain stimulation for the treatment of chronic cluster headaches to be **investigational.***

Based on review of available data, the Company considers deep brain stimulation for the treatment of other psychiatric or neurologic disorders, including but not limited to epilepsy, Tourette syndrome, depression, obsessive-compulsive disorder, anorexia nervosa, alcohol addiction, Alzheimer disease, multiple sclerosis tremor, and chronic pain, to be **investigational.***

Background/Overview

Deep Brain Stimulation

Deep brain stimulation involves the stereotactic placement of an electrode into the brain (ie, hypothalamus, thalamus, globus pallidus, or subthalamic nucleus). The electrode is initially attached to a temporary transcutaneous cable for short-term stimulation to validate treatment effectiveness. Several days later, the patient returns for permanent subcutaneous surgical implantation of the cable and a radiofrequency-coupled or battery-powered programmable stimulator. The electrode is typically implanted unilaterally on the side corresponding to the most severe symptoms. However, use of bilateral stimulation using 2 electrode arrays has also been investigated in individuals with bilateral, severe symptoms. After implantation, noninvasive programming of the neurostimulator can be adjusted to the patient's symptoms. This feature may be important for individuals with Parkinson disease, whose disease may progress over time, requiring different neurostimulation parameters. Setting the optimal neurostimulation parameters may involve the balance between optimal symptom control and appearance of adverse effects of neurostimulation, such as dysarthria, disequilibrium, or involuntary movements.

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

In 1997, the Activa^{®‡} Tremor Control System (Medtronic) was approved by the U.S. Food and Drug Administration (FDA) through the pre-market approval process for deep brain stimulation. The Activa Tremor Control System consists of an implantable neurostimulator, a deep brain stimulator

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

lead, an extension that connects the lead to the power source, a console programmer, a software cartridge to set electrical parameters for stimulation, and a patient control magnet, which allows the patient to turn the neurostimulator on and off, or change between high and low settings.

The FDA-labeled indications for Activa were originally limited to unilateral implantation for the treatment of tremor, but the indications have evolved over time. In 2002, the FDA labeled indications were expanded to include bilateral implantation as a treatment to decrease the symptoms of advanced Parkinson disease not controlled by medication. In 2003, the labeled indications were further expanded to include "...unilateral or bilateral stimulation of the internal globus pallidus or subthalamic nucleus to aid in the management of chronic, intractable (drug refractory) primary dystonia, including generalized and/or segmental dystonia, hemidystonia, and cervical dystonia (torticollis) in patients 7 years of age or above." In 2018, the deep brain stimulation system received an expanded indication as an adjunctive therapy for epilepsy (P960009-S219). Other deep brain stimulation systems are described in Table 1.

System	Manufacturer	FDA Product Code	PMA or HDE	Approval Date	Indications
Activa ^{®‡} Deep Brain Stimulation Therapy System	Medtronic	MBX	P96009	1997	Unilateral or bilateral stimulation of the internal globus pallidus or subthalamic nucleus for symptoms of Parkinson disease or primary dystonia
Reclaim ^{®‡} DBS Therapy for Obsessive Compulsive Disorder	Medtronic		H050003	2009	Bilateral stimulation of the anterior limb of the internal capsule for severe obsessive-compulsive disorder

 Table 1. Deep Brain Stimulation Systems

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

Brio Neurostimulation System	St. Jude Medical	NHL	P140009	2015	Parkinsonian tremor (subthalamic nucleus) and essential tremor (thalamus)
Infinity DBS	Abbott Medical/St. Jude Medical	PJS	P140009	2016	Parkinsonian tremor
Vercise DBS System	Boston Scientific	NHL	P150031	2017	Moderate-to-advanced levodopa-responsive PD inadequately controlled with medication alone
Medtronic DBS System for Epilepsy	Medtronic	MBX	P960009- S219	2018	Expanded indication for epilepsy with bilateral stimulation of the anterior nucleus of the thalamus
Percept PC Deep Brain Stimulation	Medtronic	MHY	P960009- S	2020	Records brain signals while delivering therapy for PD or primary dystonia
Vercise Genus DBS System	Boston Scientific	NHL	P150031- S034	2021	Stimulation of the subthalamic nucleus and globus pallidus for PD
SenSight Directional Lead System	Medtronic	МНҮ	P960009	2021	Unilateral or bilateral stimulation for PD, tremor, dystonia, and epilepsy

DBS: deep brain stimulation; HDE: humanitarian device exemption; PD: Parkinson disease; PMA: premarket approval

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

practice in this community, technology evaluation centers, reference to federal regulations, other plan medical policies, and accredited national guidelines.

Deep brain stimulation involves the stereotactic placement of an electrode into a central nervous system nucleus (eg, hypothalamus, thalamus, globus pallidus, subthalamic nucleus). Deep brain stimulation is used as an alternative to permanent neuroablative procedures for control of essential tremor and Parkinson disease. Deep brain stimulation is also being evaluated for the treatment of a variety of other neurologic and psychiatric disorders.

Summary of Evidence

For individuals who have essential tremor or tremor in Parkinson disease who receive deep brain stimulation of the thalamus, the evidence includes a systematic review and case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The systematic review (a TEC Assessment) concluded that there was sufficient evidence that deep brain stimulation of the thalamus results in clinically significant tremor suppression and that outcomes after deep brain stimulation were at least as good as thalamotomy. Subsequent studies reporting long-term follow-up have supported the conclusions of the TEC Assessment and found that tremors were effectively controlled 5 to 6 years after deep brain stimulation. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have symptoms (eg, speech, motor fluctuations) associated with Parkinson disease (advanced or >4 years in duration with early motor symptoms) who receive deep brain stimulation of the globus pallidus interna or subthalamic nucleus, the evidence includes randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. One of the systematic reviews (a TEC Assessment) concluded that studies evaluating deep brain stimulation of the globus pallidus interna or subthalamic nucleus have consistently demonstrated clinically significant improvements in outcomes (eg, neurologic function). Other systematic reviews have also found significantly better outcomes after deep brain stimulation than after a control intervention. An RCT in patients with levodopa-responsive Parkinson disease of at least 4 years in duration and uncontrolled motor symptoms found that quality of life at 2 years was significantly higher when deep brain stimulation was provided in addition to medical therapy. Meta-analyses of RCTs comparing deep brain stimulation of the globus pallidus interna with deep brain stimulation is superior to the other.

^{©2024} Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have primary dystonia who receive deep brain stimulation of the globus pallidus interna or subthalamic nucleus, the evidence includes systematic reviews, RCTs, and case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. A pooled analysis of 24 studies, mainly uncontrolled, found improvements in motor scores and disability scores after 6 months and at last follow-up (mean, 32 months). Both double-blind RCTs found that severity scores improved more after active than after sham stimulation. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have tardive dyskinesia or tardive dystonia who receive deep brain stimulation, the evidence includes an RCT and case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The RCT did not report statistically significant improvement in the dystonia severity outcomes or the secondary outcomes related to disability and quality of life, but these may have been underpowered. Additional studies, especially RCTs or other controlled studies, are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have epilepsy who receive deep brain stimulation, the evidence includes systematic reviews, RCTs, and many observational studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Two RCTs with more than 15 patients were identified. The first RCT (N=110) evaluated anterior thalamic nucleus deep brain stimulation and reported that deep brain stimulation had a positive impact on seizure frequency during some parts of the blinded trial phase, but not others, and a substantial number of adverse events (in >30% of patients). There were no differences between groups in 50% responder rates, Liverpool Seizure Severity Scale, or Quality of Life in Epilepsy scores. A 7-year open-label follow-up of the RCT included 66% of implanted patients; reasons for missing data were primarily related to adverse events or dissatisfaction with the device. Reduction in seizure frequency continued to improve during follow-up among the patients who continued follow-up. The second RCT (N=16) showed a benefit with deep brain stimulation. Many observational studies reported fewer seizures compared with baseline, however, without control groups, interpretation of these results is limited. Additional trials are required to determine the impact of deep brain stimulation on patient outcomes.

^{©2024} Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have Tourette syndrome who receive deep brain stimulation, the evidence includes observational studies, RCTs, and systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Two RCTs with 15 or more patients have been reported. One RCT found differences in severity of Tourette syndrome for active versus sham at 3 months while the other RCT did not. Neither study demonstrated improvements in comorbid symptoms of obsessive-compulsive disorder or depression. Both studies reported high rates of serious adverse events. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have cluster headaches or facial pain who receive deep brain stimulation, the evidence includes a systematic review, randomized crossover study, and case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The systematic review included an individual patient data meta-analysis of 34 patients, showing a significant reduction in pain intensity at 3 months following deep brain stimulation for chronic facial pain; data for follow-up beyond 3 months were not eligible for statistical analysis. In an RCT of 11 patients with severe, refractory, chronic cluster headache, the between-group difference in response rates did not differ significantly between active and sham stimulation phases. Additional RCTs or controlled studies are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have treatment-resistant depression who receive deep brain stimulation, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. A number of case series and several prospective controlled trials evaluating deep brain stimulation have been published. Two RCTs of deep brain stimulation in the subgenual cingulate cortex and ventral striatum/ventral capsule were terminated for futility. Another RCT of stimulation of the same brain area (ventral striatum/ventral capsule) did not find a statistically significant difference between groups in the primary outcome (clinical response), and adverse psychiatric events occurred more frequently in the treatment group than in the control group. More recently, a controlled crossover trial randomized patients to sham or active stimulation of the anterior limb of the internal capsule after a year of open-label stimulation. There was a greater reduction in symptom scores after active stimulation, but only in patients who

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

were responders in the open-label phase. Stimulation of the subcallosal (subgenual) cingulate was evaluated in a 2019 sham-controlled within-subject study that found prolonged response in 50% of patients and remission in 30% of patients with treatment-resistant depression. Deep brain stimulation for patients with major depressive disorder who have failed all other treatment options is an active area of research, but the brain regions that might prove to be effective for treatment-resistant depression have yet to be established. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have obsessive-compulsive disorder who receive deep brain stimulation, the evidence includes meta-analyses of RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Among the RCTs on deep brain stimulation for obsessive-compulsive disorder included in meta-analyses, only 1 has reported an outcome of clinical interest (therapeutic response rate), and that trial did not find a statistically significant benefit for deep brain stimulation compared with sham treatment. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have other neurologic or psychiatric disorders who receive deep brain stimulation, the evidence includes a number of nonrandomized studies or RCTs in patients with multiple sclerosis, chronic pain, or alcohol use disorder. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. One RCT with 10 multiple sclerosis patients, 2 RCTs in patients with chronic pain, and 1 RCT in patients with treatment-refractory alcohol use disorder is insufficient evidence on which to draw conclusions about the efficacy of deep brain stimulation in these populations. Additional trials are required. For individuals who have anorexia nervosa, Alzheimer disease, Huntington disease, or chronic pain who receive deep brain stimulation for these conditions. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers,

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2014 Input

In response to requests, input was received from 2 academic medical centers and 2 physician specialty societies while this policy was under review in 2014. Input supported the use of bilateral deep brain stimulation in individuals with medically unresponsive tremor in both limbs.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Academy of Neurology

Essential Tremor

In 2011, the American Academy of Neurology (AAN) updated its guidelines on the treatment of essential tremor, which were reaffirmed in 2022. This update did not change the conclusions and recommendations of the AAN (2005) practice parameters on deep brain stimulation for essential tumor. The guidelines stated that bilateral deep brain stimulation of the thalamic nucleus may be used to treat medically refractory limb tremor in both upper limbs (level C, possibly effective) but that there were insufficient data on the risk/benefit ratio of bilateral versus unilateral deep brain stimulation in the treatment of limb tremor. There was insufficient evidence to make recommendations on the use of thalamic deep brain stimulation for head or voice tremor (level U, treatment is unproven).

Parkinson Disease

In 2018, the AAN affirmed the guideline developed by the Congress of Neurological Surgeons (see Table 2).

Tourette Syndrome

Guidelines from AAN (2019, reaffirmed 2022) provide recommendations on the assessment for and use of deep brain stimulation in adults with severe, treatment-refractory tics. The AAN notes that

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

patients with severe Tourette syndrome resistant to medical and behavioral therapy may benefit from deep brain stimulation, but there is no consensus on the optimal brain target. Brain regions that have been stimulated in patients with Tourette syndrome include the centromedian thalamus, the globus pallidus internus (ventral and dorsal), the globus pallidus externus, the subthalamic nucleus, and the ventral striatum/ventral capsular nucleus accumbens region. The AAN concludes that deep brain stimulation of the anteromedial globus pallidus is possibly more likely than sham stimulation to reduce tic severity.

American Society for Stereotactic and Functional Neurosurgery

Obsessive-Compulsive Disorder

In 2021, the American Society for Stereotactic and Functional Neurosurgery and the Congress of Neurological Surgeons updated their 2014 guidelines on deep brain stimulation for obsessive-compulsive disorder. The document concluded that there was a single level I study supporting the use of bilateral subthalamic nucleus deep brain stimulation for medically refractory obsessive-compulsive disorder and a single level II study supporting bilateral nucleus accumbens or bed nucleus of stria terminalis deep brain stimulation for medically refractory obsessive-compulsive disorder. It also concluded that the evidence on unilateral deep brain stimulation was insufficient.

Refractory Epilepsy

In 2022, the American Society for Stereotactic and Functional Neurosurgery published a position statement on deep brain stimulation for medication-refractory epilepsy. Indications for deep brain stimulation include confirmed diagnosis of epilepsy (focal onset seizures with or without generalization), failure to achieve seizure control after 2 or more appropriately dosed seizure medications, seizures with localized onset in a region that cannot be resected or for which surgical resection has failed, or focal-onset seizures with a nonlocalized or unclear region of onset.

Congress of Neurologic Surgeons Parkinson Disease

In 2018, evidence-based guidelines from the Congress of Neurologic Surgeons, affirmed by the AAN, compared the efficacy of bi-lateral deep brain stimulation of the subthalamic nucleus and globus pallidus internus for the treatment of patients with Parkinson disease.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

 Table 2. Recommendations of the Congress of Neurologic Surgeons for DBS for Parkinson

 Disease

Goal	Most Effective Area of Stimulation (subthalamic nucleus or globus pallidus internus)	Level of Evidence
Improving motor symptoms	subthalamic nucleus or globus pallidus internus are similarly effective	Ι
Reduction of dopaminergic medication	subthalamic nucleus	Ι
Treatment of "on" medication dyskinesias	globus pallidus internus if reduction of medication is not anticipated	Ι
Quality of life	no evidence to recommend one over the other	Ι
Lessen impact of DBS on cognitive decline	globus pallidus internus	Ι
Reduce risk of depression	globus pallidus internus	Ι
Reduce adverse effects	insufficient evidence to recommend one over the other	Insufficient

DBS: Deep brain stimulation

National Institute for Health and Care Excellence

The United Kingdom's NICE has published guidance documents on deep brain stimulation, as discussed in the following subsections.

Tremor and Dystonia

In 2006, NICE made the same statements about use of deep brain stimulation for treatment of both tremor and dystonia. Unilateral and bilateral stimulation of structures responsible for modifying movements, such as the thalamus, globus pallidus, and the subthalamic nucleus, which interact functionally with the substantia nigra, are included in both guidance statements. The guidance stated:

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

"Current evidence on the safety and efficacy of deep brain stimulation for tremor and dystonia (excluding Parkinson's disease) appears adequate to support the use of this procedure."

Refractory Chronic Pain Syndromes (Excluding Headache)

In 2011, guidance from NICE indicated there is evidence that deep brain stimulation for refractory chronic pain (excluding headache) is associated with serious risks. However, the procedure is "efficacious in some patients" refractory to other treatments." Patients should be informed that deep brain stimulation may not control their chronic pain symptoms and that possible risks associated with this procedure include the small risk of death.

Intractable Trigeminal Autonomic Cephalalgias

In 2011, guidance from NICE indicated that the evidence on the efficacy of deep brain stimulation for intractable trigeminal autonomic cephalalgias (eg, cluster headaches) was "limited and inconsistent, and the evidence on safety shows that there were serious but well-known adverse effects."

Refractory Epilepsy

In 2020, guidance from NICE indicated that the evidence on the efficacy and safety of deep brain stimulation for refractory epilepsy (for anterior thalamic targets) was limited in both quantity and quality, and "this procedure should only be used with special arrangements for clinical governance, consent, and audit or research". For targets other than the anterior thalamus, NICE recommends that "this procedure should only be used in the context of research".

Parkinson Disease

In 2003, NICE stated that the evidence on the safety and efficacy of deep brain stimulation for treatment of Parkinson disease "appears adequate to support the use of the procedure." The guidance noted that deep brain stimulation should only be offered when Parkinson disease is refractory to best medical treatment.

U.S. Preventive Services Task Force Recommendations

Not applicable.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

Medicare National Coverage

Effective for services furnished in April 2003, Medicare covers unilateral or bilateral thalamic ventralis intermedius nucleus deep brain stimulation for the treatment of essential tumor and/or parkinsonian tremor and unilateral or bilateral subthalamic nucleus or globus pallidus interna deep brain stimulation for the treatment of Parkinson disease when the following conditions are met:

- 1. Devices must be approved by the Food and Drug Administration (FDA) for "deep brain stimulation or devices used in accordance with FDA-approved protocols governing Category B Investigational Device Exemption (IDE) deep brain stimulation clinical trials."
- 2. For thalamic ventralis intermedius nucleus deep brain stimulation, patients must meet all of the following criteria:
 - a. "Diagnosis of ET [essential t remor] based on postural or kinetic tremors of hand(s) without other neurologic signs, or diagnosis of idiopathic Parkinson disease (presence of at least 2 cardinal PD [Parkinson disease] features (tremor, rigidity or bradykinesia)) which is of a tremor-dominant form.
 - b. Marked disabling tremor of at least level 3 or 4 on the Fahn-Tolosa-Marin Clinical Tremor Rating Scale (or equivalent scale) in the extremity intended for treatment, causing significant limitation in daily activities despite optimal medical therapy.
 - c. Willingness and ability to cooperate during conscious operative procedure, as well as during postsurgical evaluations, adjustments of medications and stimulator settings."
- 3. For subthalamic nucleus or globus pallidus interna deep brain stimulation, patients must meet all of the following criteria:
 - a. "Diagnosis of PD based on the presence of at least 2 cardinal Parkinson disease features (tremor, rigidity or bradykinesia).
 - b. Advanced idiopathic PD as determined by the use of Hoehn and Yahr stage or Unified Parkinson's Disease Rating Scale part III motor subscale.
 - c. L-dopa responsive with clearly defined 'on' periods.
 - d. Persistent disabling Parkinson's symptoms or drug side effects (eg, dyskinesias, motor fluctuations, or disabling 'off' periods) despite optimal medical therapy.
 - e. Willingness and ability to cooperate during conscious operative procedure, as well as during post-surgical evaluations, adjustments of medications and stimulator settings."

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

Deep brain stimulation is not covered for essential tumor or Parkinson disease patients with any of the following:

- 1. "Non-idiopathic Parkinson's disease or 'Parkinson's Plus' syndromes.
- 2. Cognitive impairment, dementia or depression, which would be worsened by or would interfere with the patient's ability to benefit from DBS [deep brain stimulation].
- 3. Current psychosis, alcohol abuse or other drug abuse.

Structural lesions such as basal ganglionic stroke, tumor or vascular malformation as etiology of the movement disorder.

Previous movement disorder surgery within the affected basal ganglion.

Significant medical, surgical, neurologic or orthopedic co-morbidities contraindicating DBS [deep brain stimulation] surgery or stimulation."

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 3. Included are randomized controlled trials with at least 40 participants, excluding trials on deep brain stimulation for Parkinson disease.

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
Epilepsy			
NCT04181229	Deep Brain Stimulation After Failed Vagal Nerve Stimulation for the Treatment of Drug-Resistant Epilepsy in Children	50	Mar 2025
NCT04164056	Hippocampal and Thalamic deep brain stimulation for Bilateral Temporal Lobe Epilepsy	80	Sep 2024

Table 3. Summary of Key Trials

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

NCT No.	Trial Name	Planned Enrollment	Completion Date
NCT03900468ª	Medtronic Deep Brain Stimulation Therapy for Epilepsy Post-Approval Study (EPAS)	140	Mar 2028
NCT06248333	Subthalamic Nucleus Electrical Stimulation for Drug-resistant Focal Motor Epilepsy (STEM)	33	Jan 2026
Huntington's Disease			
NCT04244513ª	Deep Brain Stimulation Treatment for Chorea in Huntington's Disease	40	Dec 2023
Obsessive- Compulsive Disorder			
NCT02773082ª	Reclaim Deep Brain Stimulation Therapy for Obsessive-Compulsive Disorder (OCD)	50	Jan 2030
NCT02844049	European Study of Quality of Life in Resistant OCD Patients Treated by subthalamic nucleus deep brain stimulation	60	Dec 2026
NCT05995951	Deep Brain Stimulation Surgery for the Treatment of Refractory Obsessive-Compulsive Disorder	10	Sept 2025
Treatment Resistant Depression			
NCT03653858ª	Controlled Randomized Clinical Trial to Assess Efficacy of Deep Brain Stimulation of the slMFB in Patients With Treatment Resistant Major Depression (FORSEEIII)	47	Jun 2025
NCT06096207	DBS for Depression	20	Oct 2028

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

NCT No.	Trial Name	Planned Enrollment	Completion Date
Alzheimer Disease			
NCT03622905	Advance II Study: DBS-f in Patients With Mild Alzheimer's Disease	210	Oct 2025
NCT05882344	Deep Brain Stimulation for Alzheimer's	8	Oct 2028
NCT05762926	Non-invasive Brain Stimulation by Transcranial Pulse Stimulation as a Coadjunctive Treatment in Alzheimer's Disease	50	May 2024
Unpublished			
NCT02076698	Deep Brain Stimulation of the Anterior Nucleus of the Thalamus in Epilepsy	62	Nov 2021

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

References

- Schuurman PR, Bosch DA, Merkus MP, et al. Long-term follow-up of thalamic stimulation versus thalamotomy for tremor suppression. Mov Disord. Jun 15 2008; 23(8): 1146-53. PMID 18442104
- Hariz MI, Krack P, Alesch F, et al. Multicentre European study of thalamic stimulation for parkinsonian tremor: a 6 year follow-up. J Neurol Neurosurg Psychiatry. Jun 2008; 79(6): 694-9. PMID 17898034
- 3. Putzke JD, Uitti RJ, Obwegeser AA, et al. Bilateral thalamic deep brain stimulation: midline tremor control. J Neurol Neurosurg Psychiatry. May 2005; 76(5): 684-90. PMID 15834027
- 4. Pahwa R, Lyons KE, Wilkinson SB, et al. Long-term evaluation of deep brain stimulation of the thalamus. J Neurosurg. Apr 2006; 104(4): 506-12. PMID 16619653
- Jost ST, Aloui S, Evans J, et al. Neurostimulation for Advanced Parkinson Disease and Quality of Life at 5 Years: A Nonrandomized Controlled Trial. JAMA Netw Open. Jan 02 2024; 7(1): e2352177. PMID 38236600

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

- 6. Schnitzler A, Mir P, Brodsky MA, et al. Directional Deep Brain Stimulation for Parkinson's Disease: Results of an International Crossover Study With Randomized, Double-Blind Primary Endpoint. Neuromodulation. Aug 2022; 25(6): 817-828. PMID 34047410
- Perestelo-Pérez L, Rivero-Santana A, Pérez-Ramos J, et al. Deep brain stimulation in Parkinson's disease: meta-analysis of randomized controlled trials. J Neurol. Nov 2014; 261(11): 2051-60. PMID 24487826
- 8. Kleiner-Fisman G, Herzog J, Fisman DN, et al. Subthalamic nucleus deep brain stimulation: summary and meta-analysis of outcomes. Mov Disord. Jun 2006; 21 Suppl 14: S290-304. PMID 16892449
- 9. Appleby BS, Duggan PS, Regenberg A, et al. Psychiatric and neuropsychiatric adverse events associated with deep brain stimulation: A meta-analysis of ten years' experience. Mov Disord. Sep 15 2007; 22(12): 1722-8. PMID 17721929
- 10. Schuepbach WM, Rau J, Knudsen K, et al. Neurostimulation for Parkinson's disease with early motor complications. N Engl J Med. Feb 14 2013; 368(7): 610-22. PMID 23406026
- 11. Sako W, Miyazaki Y, Izumi Y, et al. Which target is best for patients with Parkinson's disease? A meta-analysis of pallidal and subthalamic stimulation. J Neurol Neurosurg Psychiatry. Sep 2014; 85(9): 982-6. PMID 24444854
- 12. Combs HL, Folley BS, Berry DT, et al. Cognition and Depression Following Deep Brain Stimulation of the Subthalamic Nucleus and Globus Pallidus Pars Internus in Parkinson's Disease: A Meta-Analysis. Neuropsychol Rev. Dec 2015; 25(4): 439-54. PMID 26459361
- 13. Tan ZG, Zhou Q, Huang T, et al. Efficacies of globus pallidus stimulation and subthalamic nucleus stimulation for advanced Parkinson's disease: a meta-analysis of randomized controlled trials. Clin Interv Aging. 2016; 11: 777-86. PMID 27382262
- 14. Wang JW, Zhang YQ, Zhang XH, et al. Cognitive and Psychiatric Effects of STN versus GPi Deep Brain Stimulation in Parkinson's Disease: A Meta-Analysis of Randomized Controlled Trials. PLoS One. 2016; 11(6): e0156721. PMID 27248139
- 15. Xie CL, Shao B, Chen J, et al. Effects of neurostimulation for advanced Parkinson's disease patients on motor symptoms: A multiple-treatments meta-analysas of randomized controlled trials. Sci Rep. May 04 2016; 6: 25285. PMID 27142183
- 16. Xu F, Ma W, Huang Y, et al. Deep brain stimulation of pallidal versus subthalamic for patients with Parkinson's disease: a meta-analysis of controlled clinical trials. Neuropsychiatr Dis Treat. 2016; 12: 1435-44. PMID 27382286

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

- 17. Wong JK, Cauraugh JH, Ho KWD, et al. STN vs. GPi deep brain stimulation for tremor suppression in Parkinson disease: A systematic review and meta-analysis. Parkinsonism Relat Disord. Jan 2019; 58: 56-62. PMID 30177491
- 18. U.S. Food and Drug Administration. Summary of Safety and Probable Benefit. Medtronic Activa Dystonia Therapy. 2003; http://www.accessdata.fda.gov/cdrh_docs/pdf2/H020007b.pdf.
- 19. Moro E, LeReun C, Krauss JK, et al. Efficacy of pallidal stimulation in isolated dystonia: a systematic review and meta-analysis. Eur J Neurol. Apr 2017; 24(4): 552-560. PMID 28186378
- 20. Rodrigues FB, Duarte GS, Prescott D, et al. Deep brain stimulation for dystonia. Cochrane Database Syst Rev. Jan 10 2019; 1(1): CD012405. PMID 30629283
- 21. Kupsch A, Benecke R, Müller J, et al. Pallidal deep-brain stimulation in primary generalized or segmental dystonia. N Engl J Med. Nov 09 2006; 355(19): 1978-90. PMID 17093249
- 22. Volkmann J, Mueller J, Deuschl G, et al. Pallidal neurostimulation in patients with medicationrefractory cervical dystonia: a randomised, sham-controlled trial. Lancet Neurol. Sep 2014; 13(9): 875-84. PMID 25127231
- Gruber D, Südmeyer M, Deuschl G, et al. Neurostimulation in tardive dystonia/dyskinesia: A delayed start, sham stimulation-controlled randomized trial. Brain Stimul. 2018; 11(6): 1368-1377. PMID 30249417
- 24. Damier P, Thobois S, Witjas T, et al. Bilateral deep brain stimulation of the globus pallidus to treat tardive dyskinesia. Arch Gen Psychiatry. Feb 2007; 64(2): 170-6. PMID 17283284
- 25. Pouclet-Courtemanche H, Rouaud T, Thobois S, et al. Long-term efficacy and tolerability of bilateral pallidal stimulation to treat tardive dyskinesia. Neurology. Feb 16 2016; 86(7): 651-9. PMID 26791148
- 26. Kwan P, Arzimanoglou A, Berg AT, et al. Definition of drug resistant epilepsy: consensus proposal by the ad hoc Task Force of the ILAE Commission on Therapeutic Strategies. Epilepsia. Jun 2010; 51(6): 1069-77. PMID 19889013
- 27. Borghs S, de la Loge C, Cramer JA. Defining minimally important change in QOLIE-31 scores: estimates from three placebo-controlled lacosamide trials in patients with partial-onset seizures. Epilepsy Behav. Mar 2012; 23(3): 230-4. PMID 22341962
- 28. Sprengers M, Vonck K, Carrette E, et al. Deep brain and cortical stimulation for epilepsy. Cochrane Database Syst Rev. Jul 18 2017; 7(7): CD008497. PMID 28718878
- 29. Li MCH, Cook MJ. Deep brain stimulation for drug-resistant epilepsy. Epilepsia. Feb 2018; 59(2): 273-290. PMID 29218702

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

- Bouwens van der Vlis TAM, Schijns OEMG, Schaper FLWVJ, et al. Deep brain stimulation of the anterior nucleus of the thalamus for drug-resistant epilepsy. Neurosurg Rev. Jun 2019; 42(2): 287-296. PMID 29306976
- 31. Fisher R, Salanova V, Witt T, et al. Electrical stimulation of the anterior nucleus of thalamus for treatment of refractory epilepsy. Epilepsia. May 2010; 51(5): 899-908. PMID 20331461
- 32. Food and Drug Administration. Medtronic DBS System for Epilepsy, Summary of Safety and Effectiveness Data (SSED).
- 33. Tröster AI, Meador KJ, Irwin CP, et al. Memory and mood outcomes after anterior thalamic stimulation for refractory partial epilepsy. Seizure. Feb 2017; 45: 133-141. PMID 28061418
- 34. Cukiert A, Cukiert CM, Burattini JA, et al. Seizure outcome after hippocampal deep brain stimulation in patients with refractory temporal lobe epilepsy: A prospective, controlled, randomized, double-blind study. Epilepsia. Oct 2017; 58(10): 1728-1733. PMID 28744855
- 35. Salanova V, Witt T, Worth R, et al. Long-term efficacy and safety of thalamic stimulation for drug-resistant partial epilepsy. Neurology. Mar 10 2015; 84(10): 1017-25. PMID 25663221
- 36. Kim SH, Lim SC, Kim J, et al. Long-term follow-up of anterior thalamic deep brain stimulation in epilepsy: A 11-year, single center experience. Seizure. Nov 2017; 52: 154-161. PMID 29040867
- 37. Peltola J, Colon AJ, Pimentel J, et al. Deep Brain Stimulation of the Anterior Nucleus of the Thalamus in Drug-Resistant Epilepsy in the MORE Multicenter Patient Registry. Neurology. May 02 2023; 100(18): e1852-e1865. PMID 36927882
- 38. Yan H, Wang X, Zhang X, et al. Deep brain stimulation for patients with refractory epilepsy: nuclei selection and surgical outcome. Front Neurol. 2023; 14: 1169105. PMID 37251216
- 39. Baldermann JC, Schüller T, Huys D, et al. Deep Brain Stimulation for Tourette-Syndrome: A Systematic Review and Meta-Analysis. Brain Stimul. 2016; 9(2): 296-304. PMID 26827109
- 40. Fraint A, Pal G. Deep Brain Stimulation in Tourette's Syndrome. Front Neurol. 2015; 6: 170. PMID 26300844
- 41. Schrock LE, Mink JW, Woods DW, et al. Tourette syndrome deep brain stimulation: a review and updated recommendations. Mov Disord. Apr 2015; 30(4): 448-71. PMID 25476818
- 42. Servello D, Zekaj E, Saleh C, et al. Sixteen years of deep brain stimulation in Tourette's Syndrome: a critical review. J Neurosurg Sci. Jun 2016; 60(2): 218-29. PMID 26788742
- 43. Piedad JC, Rickards HE, Cavanna AE. What patients with gilles de la tourette syndrome should be treated with deep brain stimulation and what is the best target?. Neurosurgery. Jul 2012; 71(1): 173-92. PMID 22407075

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

- 44. Wehmeyer L, Schüller T, Kiess J, et al. Target-Specific Effects of Deep Brain Stimulation for Tourette Syndrome: A Systematic Review and Meta-Analysis. Front Neurol. 2021; 12: 769275. PMID 34744993
- 45. Kefalopoulou Z, Zrinzo L, Jahanshahi M, et al. Bilateral globus pallidus stimulation for severe Tourette's syndrome: a double-blind, randomised crossover trial. Lancet Neurol. Jun 2015; 14(6): 595-605. PMID 25882029
- 46. Welter ML, Houeto JL, Thobois S, et al. Anterior pallidal deep brain stimulation for Tourette's syndrome: a randomised, double-blind, controlled trial. Lancet Neurol. Aug 2017; 16(8): 610-619. PMID 28645853
- 47. Martinez-Ramirez D, Jimenez-Shahed J, Leckman JF, et al. Efficacy and Safety of Deep Brain Stimulation in Tourette Syndrome: The International Tourette Syndrome Deep Brain Stimulation Public Database and Registry. JAMA Neurol. Mar 01 2018; 75(3): 353-359. PMID 29340590
- 48. International Headache Society. International Classification of Headache Disorders. 2018; https://www.ichd-3.org.
- 49. Qassim H, Zhao Y, Ströbel A, et al. Deep Brain Stimulation for Chronic Facial Pain: An Individual Participant Data (IPD) Meta-Analysis. Brain Sci. Mar 14 2023; 13(3). PMID 36979302
- 50. Fontaine D, Lazorthes Y, Mertens P, et al. Safety and efficacy of deep brain stimulation in refractory cluster headache: a randomized placebo-controlled double-blind trial followed by a 1-year open extension. J Headache Pain. Feb 2010; 11(1): 23-31. PMID 19936616
- 51. Bussone G, Franzini A, Proietti Cecchini A, et al. Deep brain stimulation in craniofacial pain: seven years' experience. Neurol Sci. May 2007; 28 Suppl 2: S146-9. PMID 17508162
- 52. Broggi G, Franzini A, Leone M, et al. Update on neurosurgical treatment of chronic trigeminal autonomic cephalalgias and atypical facial pain with deep brain stimulation of posterior hypothalamus: results and comments. Neurol Sci. May 2007; 28 Suppl 2: S138-45. PMID 17508161
- 53. Mandat V, Zdunek PR, Krolicki B, et al. Periaqueductal/periventricular gray deep brain stimulation for the treatment of neuropathic facial pain. Front Neurol. 2023; 14: 1239092. PMID 38020618
- 54. Sobstyl M, Kupryjaniuk A, Prokopienko M, et al. Subcallosal Cingulate Cortex Deep Brain Stimulation for Treatment-Resistant Depression: A Systematic Review. Front Neurol. 2022; 13: 780481. PMID 35432155

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

- 55. Hitti FL, Yang AI, Cristancho MA, et al. Deep Brain Stimulation Is Effective for Treatment-Resistant Depression: A Meta-Analysis and Meta-Regression. J Clin Med. Aug 30 2020; 9(9). PMID 32872572
- 56. Wu Y, Mo J, Sui L, et al. Deep Brain Stimulation in Treatment-Resistant Depression: A Systematic Review and Meta-Analysis on Efficacy and Safety. Front Neurosci. 2021; 15: 655412. PMID 33867929
- 57. Dougherty DD, Rezai AR, Carpenter LL, et al. A Randomized Sham-Controlled Trial of Deep Brain Stimulation of the Ventral Capsule/Ventral Striatum for Chronic Treatment-Resistant Depression. Biol Psychiatry. Aug 15 2015; 78(4): 240-8. PMID 25726497
- 58. Bergfeld IO, Mantione M, Hoogendoorn ML, et al. Deep Brain Stimulation of the Ventral Anterior Limb of the Internal Capsule for Treatment-Resistant Depression: A Randomized Clinical Trial. JAMA Psychiatry. May 01 2016; 73(5): 456-64. PMID 27049915
- 59. Crowell AL, Riva-Posse P, Holtzheimer PE, et al. Long-Term Outcomes of Subcallosal Cingulate Deep Brain Stimulation for Treatment-Resistant Depression. Am J Psychiatry. Nov 01 2019; 176(11): 949-956. PMID 31581800
- 60. Gadot R, Najera R, Hirani S, et al. Efficacy of deep brain stimulation for treatment-resistant obsessive-compulsive disorder: systematic review and meta-analysis. J Neurol Neurosurg Psychiatry. Sep 20 2022. PMID 36127157
- 61. Mar-Barrutia L, Real E, Segalás C, et al. Deep brain stimulation for obsessive-compulsive disorder: A systematic review of worldwide experience after 20 years. World J Psychiatry. Sep 19 2021; 11(9): 659-680. PMID 34631467
- 62. Raviv N, Staudt MD, Rock AK, et al. A Systematic Review of Deep Brain Stimulation Targets for Obsessive Compulsive Disorder. Neurosurgery. Nov 16 2020; 87(6): 1098-1110. PMID 32615588
- 63. Kisely S, Hall K, Siskind D, et al. Deep brain stimulation for obsessive-compulsive disorder: a systematic review and meta-analysis. Psychol Med. Dec 2014; 44(16): 3533-42. PMID 25066053
- 64. Brandmeir NJ, Murray A, Cheyuo C, et al. Deep Brain Stimulation for Multiple Sclerosis Tremor: A Meta-Analysis. Neuromodulation. Jun 2020; 23(4): 463-468. PMID 31755637
- 65. Chagot C, Bustuchina Vlaicu M, Frismand S, et al. Deep brain stimulation in multiple sclerosisassociated tremor. A large, retrospective, longitudinal open label study, with long-term followup. Mult Scler Relat Disord. Nov 2023; 79: 104928. PMID 37657308

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

- 66. Deer TR, Falowski S, Arle JE, et al. A Systematic Literature Review of Brain Neurostimulation Therapies for the Treatment of Pain. Pain Med. Nov 07 2020; 21(7): 1415-1420. PMID 32034418
- 67. Bach P, Luderer M, Müller UJ, et al. Deep brain stimulation of the nucleus accumbens in treatment-resistant alcohol use disorder: a double-blind randomized controlled multi-center trial. Transl Psychiatry. Feb 08 2023; 13(1): 49. PMID 36755017
- Gratwicke J, Zrinzo L, Kahan J, et al. Bilateral nucleus basalis of Meynert deep brain stimulation for dementia with Lewy bodies: A randomised clinical trial. Brain Stimul. 2020; 13(4): 1031-1039. PMID 32334074
- 69. Shaffer A, Naik A, Bederson M, et al. Efficacy of deep brain stimulation for the treatment of anorexia nervosa: a systematic review and network meta-analysis of patient-level data. Neurosurg Focus. Feb 2023; 54(2): E5. PMID 36724522
- 70. Zesiewicz TA, Elble RJ, Louis ED, et al. Evidence-based guideline update: treatment of essential tremor: report of the Quality Standards subcommittee of the American Academy of Neurology. Neurology. Nov 08 2011; 77(19): 1752-5. PMID 22013182
- 71. Zesiewicz TA, Elble R, Louis ED, et al. Practice parameter: therapies for essential tremor: report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurology. Jun 28 2005; 64(12): 2008-20. PMID 15972843
- 72. Rughani A, Schwalb JM, Sidiropoulos C, et al. Congress of Neurological Surgeons Systematic Review and Evidence-Based Guideline on Subthalamic Nucleus and Globus Pallidus Internus Deep Brain Stimulation for the Treatment of Patients With Parkinson's Disease: Executive Summary. Neurosurgery. Jun 01 2018; 82(6): 753-756. PMID 29538685
- 73. Pringsheim T, Okun MS, Müller-Vahl K, et al. Practice guideline recommendations summary: Treatment of tics in people with Tourette syndrome and chronic tic disorders. Neurology. May 07 2019; 92(19): 896-906. PMID 31061208
- 74. Staudt MD, Pouratian N, Miller JP, et al. Congress of Neurological Surgeons Systematic Review and Evidence-Based Guidelines for Deep Brain Stimulations for Obsessive-Compulsive Disorder: Update of the 2014 Guidelines. Neurosurgery. Mar 15 2021; 88(4): 710-712. PMID 33559678
- 75. Gummadavelli A, Englot DJ, Schwalb JM, et al. ASSFN Position Statement on Deep Brain Stimulation for Medication-Refractory Epilepsy. Neurosurgery. May 01 2022; 90(5): 636-641. PMID 35271523

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

- 76. National Institute for Health and Care Excellence (NICE). Deep brain stimulation for tremor and dystonia (excluding Parkinson's disease) [IPG188]. 2006; https://www.nice.org.uk/guidance/ipg188.
- 77. National Institute for Health and Care Excellence (NICE). Deep brain stimulation for refractory chronic pain syndromes (excluding headache) [IPG382]. 2011; http://guidance.nice.org.uk/IPG382.
- 78. National Institute for Health and Care Excellence (NICE). Deep brain stimulation for intractable trigeminal autonomic cephalalgias [IPG381]. 2011; http://www.nice.org.uk/IPG381.
- 79. National Institute for Health and Care Excellence (NICE). Deep brain stimulation for refractory epilepsy [IPG416]. 2020; https://www.nice.org.uk/guidance/IPG678/chapter/1-Recommendations.
- 80. National Institute for Health and Care Excellence (NICE). Deep brain stimulation for Parkinson's disease [IPG19]. 2003; https://www.nice.org.uk/guidance/ipg19.
- 81. Centers for Medicare & Medicaid (CMS). National Coverage Determination (NCD) for Deep Brain Stimulation for Essential Tremor and Parkinson's Disease (160.24). 2003; https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=279&ncdver=1&DocID=160.24&bc=gAAAABAAAAA&.

Policy History

Origi	nal Effecti	ve Date: 08/25/2005
Curre	nt Effectiv	ve Date: 10/14/2024
03/21	/2002	Medical Policy Committee review
03/25	/2002	Managed Care Advisory Council approval
06/24	/2002	Format revision. No substance change to policy.
08/03	/2004	Medical Director review
08/17	/2004	Medical Policy Committee review
08/30	/2004	Managed Care Advisory Council approval
07/14	/2005	Medical Director review
07/19	/2005	Medical Policy Committee review. Clinical criteria revision. Coverage eligibility
		changes. Added investigational statement for DBS for cluster headaches.
08/24	/2005	Managed Care Advisory Council approval
06/07	/2006	Medical Director review

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 000 Original Effect Current Effecti	ive Date: 08/25/2005
06/21/2006	Medical Policy Committee approval. Format revisions, FDA/Governmental, Rationale/Source. Coverage eligibility unchanged.
08/01/2007	Medical Director review
08/15/2007	Medical Policy Committee approval. No change to coverage eligibility.
08/06/2008	Medical Director review
08/20/2008	Medical Policy Committee approval. Tardive dyskinesia, Tourette syndrome,
	depression and epilepsy were added to the list of investigational indications.
08/06/2009	Medical Policy Committee approval
08/26/2009	Medical Policy Implementation Committee approval. Coverage eligibility
	unchanged.
07/01/2010	Medical Policy Committee approval
07/21/2010	Medical Policy Implementation Committee approval. Coverage eligibility
07/07/2011	unchanged. Madiaal Baliay Committee approval
07/20/2011	Medical Policy Committee approval Medical Policy Implementation Committee approval. Coverage eligibility
07/20/2011	unchanged.
06/28/2012	Medical Policy Committee approval
07/27/2012	Medical Policy Implementation Committee approval. Coverage eligibility
01/21/2012	unchanged.
02/04/2013	Coding Updated
06/27/2013	Medical Policy Committee approval
07/17/2013	Medical Policy Implementation Committee approval. Coverage eligibility
	unchanged.
07/10/2014	Medical Policy Committee approval
07/16/2014	Medical Policy Implementation Committee approval. Added anorexia nervosa,
	alcohol addiction, and chronic pain as investigational indications
01/01/2015	Coding Updated
08/03/2015	Coding update: ICD10 Diagnosis code section added; ICD9 Procedure code section
00/02/2015	removed.
09/03/2015	Medical Policy Committee approval Medical Policy Implementation Committee approval Added aligibility statement
09/23/2015	Medical Policy Implementation Committee approval. Added eligibility statement for bilateral DBS of thalamus for bilateral tremors and added Alzheimer disease to
	list of investigational indications
01/01/2016	Coding update
01/01/2010	e e ano

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 0002			
Original Effecti			
Current Effectiv	ve Date: 10/14/2024		
09/08/2016	Medical Policy Committee approval		
09/21/2016	Medical Policy Implementation Committee approval. Added "upper" to coverage		
	statement for bilateral DBS.		
01/01/2017	Coding update: Removing ICD-9 Diagnosis Codes		
09/07/2017	Medical Policy Committee approval		
09/20/2017	Medical Policy Implementation Committee approval. In medically necessary		
	statement on unilateral or bilateral deep brain stimulation of the globus pallidus or		
	subthalamic nucleus, "OR Parkinson disease for at least 4 years" added to medically		
	necessary criteria for use in Parkinson disease.		
09/06/2018	Medical Policy Committee approval		
09/19/2018	Medical Policy Implementation Committee approval. No change to coverage.		
09/05/2019	Medical Policy Committee approval		
09/11/2019	Medical Policy Implementation Committee approval. No change to coverage.		
09/03/2020	Medical Policy Committee approval		
09/09/2020	Medical Policy Implementation Committee approval. No change to coverage.		
12/11/2020	Coding update		
09/02/2021	Medical Policy Committee approval		
09/08/2021	Medical Policy Implementation Committee approval. No change to coverage.		
09/01/2022	Medical Policy Committee approval		
09/14/2022	Medical Policy Implementation Committee approval. No change to coverage.		
09/07/2023	Medical Policy Committee approval		
09/13/2023	Medical Policy Implementation Committee approval. No change to coverage.		
09/05/2024	Medical Policy Committee approval		
09/11/2024	Medical Policy Implementation Committee approval. No change to coverage.		
Next Scheduled Review Date: 09/2025			

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology $(CPT^{\circledast})^{\ddagger}$, copyright 2023 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

 $\textcircled{\sc c}2024$ Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code		
СРТ	61850, 61860, 61863, 61864, 61867, 61868, 61880, 61885, 61886, 95970, 95983, 95984		
HCPCS	C1816, C1820, C1822, C1823, C1883, L8680, L8681, L8682, L8683, L8684, L8685, L8686, L8687, L8688, L8689		
ICD-10 Diagnosis	All related diagnoses		

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:
 - 1. Consultation with technology evaluation center(s);
 - 2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
 - 3. Reference to federal regulations.

**Medically Necessary (or "Medical Necessity") - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, "nationally accepted standards of medical practice" means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient's health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Policy # 00024 Original Effective Date: 08/25/2005 Current Effective Date: 10/14/2024

recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.